Infection Control Using Evidence Practice to ensure health and safety of patients and staff.

Objectives

- Describe what, when, and how to perform hand hygiene
- Review the fundamentals of isolation precautions and what they mean
- Describe Tuberculosis in Alaska
- Review OSHA requirements for bloodborne pathogen training
- Discuss Employee Health Services

Infection Preventionist: Rebecca Hamel, MHI, RN

Infection Preventionist: Victoria Miltersen, RN, BS, BA,


Medical Director: Dr. Megan Clancy

Program Assistant: Sue Hargis

- Contact us at: 2-4829
 - Office Hours:
 - Monday Friday
 - 8am 4:30pm
- Contact shift supervisor with off-hours concerns



PAMC Policy Manual



Hand Hygiene

Measured by

- Patient Perception via
 Press Ganey Survey
- Direct Observations

BECAUSE WE CARE ABOUT YOU, IT'S OKAY TO ASK US IF




IT ONLY TAKES A FEW SIMPLE WORDS TO ENCOURAGE THIS HEALTHY HABIT— • "EXCUSE ME, DID YOU WASH YOUR HANDS?" • "I SAW YOU WASH YOUR HANDS—THANK YOU!" Hand Hygiene



These scores reflect survey data for our patients perception of Hand Hygiene and how we compare to other hospitals accross the US.

Please wash your hands and encourage others to as well.

it's OKAY TO ASK



At or slightly better

than US ave.

Under US Average

03

04

Above US Ave.

When Does Hand Hygiene Occur?



Skin Care Basics

- Provon soap
- Purell Hand Sanitizer
- <u>No</u> home lotions or hand sanitizers

No artificial, gel or shellac nails for patient care

Keep natural nails less than 1/4 inch long






Pop Quiz!!

How long should you wash your hands?

How long should you wash your hands?

- 20-25 seconds
- Some say to sing "Happy Birthday" in your head while washing.

Standard Precautions <u>All Patients – All the Time</u>




Wash or sanitize your hands.... Before and after Patient Contact Know when to use PPE




Use correct PPE



Put PPE on correctly



Properly clean equipment





Always check manufacturer's recommendation for equipment cleaning.





3 Minutes

In addition to Standard Precautions there are Isolation Precautions...

Contact Airborne Droplet Contact Enteric




Look for and follow instructions posted on the patients door <u>before</u> entering patient room.



For patients with MRSA, or other multi-drug resistant organism

For patients with TB or a novel influenza..



For patients with Measles, Mumps or RSV..



For patients with contagious viral or bacterial diarrhea



TUBERCULOSIS....



TB is primarily spread when a person coughs, sneezes, or talks.

TB Protection



<u>Required</u> when entering room of patients placed in <u>airborne precautions</u>.

Employee Health does fit testing and training of PAPRs for all employees who may be treating patients.

Employee Health



- Immunizations (HepB, Flu, TdAP)
- TB mask fit testing and/or PAPR training
- Employee exposures or injuries (needle stick, body fluid exposure, etc.)

-<u>After Hours</u>: Report to ER -Always inform your supervisor about any incident

